

## ONLINE APPENDIX

# Peer Effects in Adolescent Mental Health

Hussain Hadah

## Web.A Tables

Table Online Appendix Table A.1: Mental Health Question

| Variable    | Question  |
|-------------|---|
| <b>S60I</b> | How often have you had a poor appetite?   |
| <b>S46B</b> | Since school started this year, how often have you had trouble paying attention in school?                      |
| <b>S60J</b> | How often have you had trouble falling asleep or staying asleep?  |
| <b>S60L</b> | How often have you had trouble relaxing?  |
| <b>S60K</b> | How often was the following true during the past week [or month]? You felt depressed.                           |
| <b>S60O</b> | How often have you felt fearful?  |
| <b>S60N</b> | How often have you cried frequently?  |
| <b>S60M</b> | How often have you been moody?  |
| <b>S62A</b> | Do you agree or disagree with the following statement? You have a lot of energy.                                |
| <b>S62N</b> | Do you agree or disagree with the following statement? You feel like you are doing everything just about right. |
| <b>S62P</b> | Do you agree or disagree with the following statement? You feel loved and wanted.                               |
| <b>S62O</b> | Do you agree or disagree with the following statement? You feel socially accepted.                              |
| <b>S62H</b> | Do you agree or disagree with the following statement? You have a lot of good qualities.                        |
| <b>S62K</b> | Do you agree or disagree with the following statement? You have a lot to be proud of.                           |
| <b>S62M</b> | Do you agree or disagree with the following statement? You like yourself just the way you are.                  |

<sup>1</sup> These are the questions used to construct the depression index. The 15 items questions roughly translates to the 20 Center for Epidemiologic Studies Depression (CES-D) scale (Radloff 1977). To construct my mental health measure, I summed the responses to the negative questions and subtracted the responses to the positive. The final mental health measure is a normalized score with a mean of 0 and a standard deviation of 1.

Table Online Appendix Table A.2: Summary Statistics of the Full Sample, Friends, and Friends of Friends

|                                  | Full<br>Sample | Friends | Friends of<br>Friends |
|----------------------------------|----------------|---------|-----------------------|
| <b>Demographic</b>               |                |         |                       |
| Age                              | 15.02          | 14.98   | 14.98                 |
| Female                           | 0.50           | 0.51    | 0.51                  |
| White                            | 0.61           | 0.62    | 0.62                  |
| Black                            | 0.19           | 0.19    | 0.19                  |
| Asian                            | 0.07           | 0.07    | 0.07                  |
| Native American                  | 0.05           | 0.05    | 0.05                  |
| Other                            | 0.10           | 0.09    | 0.09                  |
| Married (w5)                     | 0.41           | 0.43    | 0.43                  |
| <b>Educational Outcomes</b>      |                |         |                       |
| GPA                              | 2.86           | 2.88    | 2.88                  |
| Number of Friends                | 6.20           | 6.52    | 6.57                  |
| <b>Health Outcomes</b>           |                |         |                       |
| Poor Health                      | 0.07           | 0.07    | 0.07                  |
| Feel Sick                        | 0.30           | 0.30    | 0.30                  |
| Feel Tired                       | 0.55           | 0.56    | 0.56                  |
| Miss School for Health Reasons   | 0.45           | 0.44    | 0.44                  |
| Ever Saw a Therapist             | 0.63           | 0.64    | 0.64                  |
| Miss School                      | 0.45           | 0.44    | 0.44                  |
| <b>Behavioral Risk Outcomes</b>  |                |         |                       |
| Skip School                      | 0.30           | 0.29    | 0.29                  |
| <i>During the last 12 months</i> |                |         |                       |
| Smoke                            | 0.36           | 0.36    | 0.36                  |
| Get Intoxicated                  | 0.31           | 0.31    | 0.31                  |
| Freq. Drink Alcohol              | 1.69           | 1.74    | 1.75                  |
| Freq. High on Drugs              | 0.12           | 0.12    | 0.12                  |
| <i>During the last month</i>     |                |         |                       |
| Freq. marijuana (w1)             | 1.92           | 1.76    | 1.74                  |
| Freq. marijuana (w3)             | 2.66           | 2.69    | 2.71                  |
| Freq. marijuana (w4)             | 0.46           | 0.46    | 0.46                  |
| Freq. Hang Out w/ Frnds          | 2.96           | 3.04    | 3.05                  |

Table Online Appendix Table A.2: Summary Statistics of the Full Sample, Friends, and Friends of Friends (*continued*)

|   | Full<br>Sample | Friends | Friends of<br>Friends |
|---|----------------|---------|-----------------------|
| Hurt Others b/c Drunk<br>Including Unprot. Sex (w4) | 0.61           | 0.61    | 0.61                  |
| Hurt Others b/c High<br>Including Unprot. Sex (w4)  | 0.24           | 0.24    | 0.24                  |
| Hurt Others b/c Drugs<br>Including Unprot. Sex (w4) | 0.43           | 0.42    | 0.43                  |
| Paid for Sex (w4)                                   | 0.02           | 0.02    | 0.02                  |

<sup>1</sup> Data source is the Add Health survey. Questions regarding skipping school without an excuse, frequency of smoking cigarettes, getting intoxicated, drinking alcohol, and getting high on drugs were asked over a 12 month period. For example, the question on frequency of smoking was 'During the past 12 months, how often did you smoke cigarettes?'. Questions regarding missing school for health reasons and frequency of consuming marijuana, and frequency of hanging out with friends were asked over a 30 day period. For example, the question on frequency of consuming marijuana 'During the past 30 days, on how many days did you use marijuana?'



Table Online Appendix Table A.3: Spillover Effect of Peers' Mental Unwellness on Own Mental Unwellness: Trimmed

|   | Mental Unwellness<br>(1)<br>OLS<br>(Reduced Form) | Friends<br>Mental Unwellness<br>(2)<br>First Stage | Mental Unwellness<br>(3)<br>2SLS |
|---|---|--|----------------------------------|
| Friends of Friends<br>Mental Unwellness | 0.229***<br>(0.022)                               | 0.272***<br>(0.018)                                |                                  |
| Friends Mental<br>Unwellness            |   |  | 0.839***<br>(0.061)              |
| Observations                            | 41,868  | 41,868   | 41,868                           |
| First Stage F-statistic                 |   |  | 1,422.54                         |

\*  $p < 0.1$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$

<sup>1</sup> This table presents the 2SLS estimation of the regression (Equation ??) with school fixed effects. Column (1) displays the estimation of the reduced form equation using own mental unwellness as the dependent variable. Moving on to column (2), I present the results of the first stage analysis using the average friends' depression as the dependent variable. Column (3) includes the results of the instrumental variable (2SLS) estimation. The control variables include sex, race, age, parental education, parental employment, parental occupation, and number of friends.

<sup>2</sup> Students that used all of there friendship nominations were trimmed from the analysis.

<sup>3</sup> Standard errors are clustered on the school level.

<sup>4</sup> Data source is the Add Health in-school survey.

Table Online Appendix Table A.4: Spillover Effect of Peers' Mental Unwellness on Own Mental Unwellness

|   | Mental Unwellness<br>(1)<br>OLS<br>(Reduced Form) | Friends<br>Mental Unwellness<br>(2)<br>First Stage | Mental Unwellness<br>(3)<br>2SLS |
|---|---|--|----------------------------------|
| Friends of Friends<br>Mental Unwellness | 0.159***<br>(0.018)                               | 0.238***<br>(0.020)                                |                                  |
| Friends Mental<br>Unwellness            |   |  | 0.669***<br>(0.057)              |
| Observations                            | 53,508  | 53,508   | 53,508                           |
| School x Grade FE                       | X   | X  | X                                |
| First Stage F-statistic                 |   |  | 1,386.2                          |

\*  $p < 0.1$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$

<sup>1</sup> This table presents the 2SLS estimation of the regression (Equation ??) with school x grade fixed effects. Column (1) displays the estimation of the reduced form equation using own mental unwellness as the dependent variable. Moving on to column (2), I present the results of the first stage analysis using the average friends' depression as the dependent variable. Column (3) includes the results of the instrumental variable (2SLS) estimation. The control variables include sex, race, age, parental education, parental employment, parental occupation, and number of friends.

<sup>2</sup> Standard errors are clustered on the school level.

<sup>3</sup> Data source is the Add Health in-school survey.

Table Online Appendix Table A.5: Spillover Effect of Reciprocated-Friends Mental Unwellness on Own Mental Unwellness

|   | Mental Unwellness<br>(1)<br>OLS<br>(Reduced Form) | Friends<br>Mental Unwellness<br>(2)<br>First Stage | Mental Unwellness<br>(3)<br>2SLS |
|---|---|--|----------------------------------|
| Reciprocated<br>Friends of Friends<br>Mental Unwellness | 0.105***<br>(0.013)                               | 0.178***<br>(0.013)                                |                                  |
| Reciprocated<br>Friends Mental<br>Unwellness            |   |  | 0.590***<br>(0.078)              |
| Observations  | 27,848  | 27,848   | 27,848                           |
| First Stage F-statistic                                 |   |  | 685.49                           |

\*  $p < 0.1$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$

<sup>1</sup> This table presents the 2SLS estimation of the regression (Equation ??) with school fixed effects. Column (1) displays the estimation of the reduced form equation using own mental unwellness as the dependent variable. Moving on to column (2), I present the results of the first stage analysis using the average friends' depression as the dependent variable. Column (3) includes the results of the instrumental variable (2SLS) estimation. The control variables include sex, race, age, parental education, parental employment, parental occupation, and number of friends.

<sup>2</sup> Standard errors are clustered on the school level.

<sup>3</sup> Data source is the Add Health in-school survey.

Table Online Appendix Table A.6: Spillover Effect of Peers' Mental Unwellness on Own Mental Unwellness: Adding Peers' Characteristics for Contextual Effects

|   | Mental Unwellness<br>(1)<br>OLS<br>(Reduced Form) | Friends<br>Mental Unwellness<br>(2)<br>First Stage | Mental Unwellness<br>(3)<br>IV |
|---|---|--|--------------------------------|
| Friends of Friends<br>Mental Unwellness | 0.183***<br>(0.021)                               | 0.240***<br>(0.021)                                |                                |
| Friends Mental<br>Unwellness            |   |  | 0.723***<br>(0.058)            |
| Observations                            | 37,154  | 37,154   | 38,365                         |
| First Stage F-statistic                 |   |  | 1,215.62                       |

\*  $p < 0.1$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$

<sup>1</sup> This table presents the 2SLS estimation of the regression (Equation ??) with school fixed effects. Column (1) displays the estimation of the reduced form equation using own mental unwellness as the dependent variable. Moving on to Column (2), I present the results of the first stage analysis using the average friends' depression as the dependent variable. Column (3) includes the results of the instrumental variable (IV) estimation. The control variables include sex, race, age, parental education, parental employment, parental occupation, and number of friends. I also include contextual effects of friends' characteristics, such as the gender, race, age, and parental background of friends. This helps to control for the contextual effects of friends' characteristics on the mental unwellness of the respondent.

<sup>2</sup> Standard errors are clustered on the school level.

<sup>3</sup> Data source is the Add Health in-school survey.

Table Online Appendix Table A.7: Spillover Effect of Peers' Mental Unwellness on Own Mental Unwellness: With Mutual and Non-mutual Friends of Friends

|                                      | Mental Unwellness<br>(1)<br>OLS<br>(Reduced Form) | Friends Mental Unwellness<br>(2)<br>First Stage | Mental Unwellness<br>(3)<br>IV |
|--------------------------------------|---|---|--------------------------------|
| Friends Mental Unwellness            |   |   | 1.559***<br>(0.052)            |
| Friends of Friends Mental Unwellness | 0.868***<br>(0.021)                               | 0.557***<br>(0.024)                             |                                |
| Observations                         | 54,258  | 54,258  | 54,258                         |
| First Stage F-statistic              |   |   | 7,886.53                       |

\*  $p < 0.1$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$

<sup>1</sup> This table presents the 2SLS estimation of the regression (Equation ??) with school fixed effects, but this sample includes mutual and non-mutual friends of friends. Column (1) displays the estimation of the reduced form equation using own mental unwellness as the dependent variable. Moving on to column (2), I present the results of the first stage analysis using the average friends' mental unwellness as the dependent variable. Finally, column (3) includes the results of the instrumental variable (IV) estimation. The control variables include sex, race, age, parental education, parental employment, parental occupation, and number of friends.

<sup>2</sup> Standard errors are clustered on the school level.

<sup>3</sup> Data source is the Add Health in-school survey.

Table Online Appendix Table A.8: Spillover Effect of Peers' Mental Health on Own Mental Unwellness: Different Indices

|                           | Mental Unwellness<br>(1)<br>IV | Mental Unwellness<br>(2)<br>IV |
|---------------------------|--------------------------------|--------------------------------|
| Friends Mental Unwellness | 0.739***<br>(0.053)            | 0.706***<br>(0.139)            |
| Observations              | 53,725                         | 6,434                          |
| School FE                 |                                | X                              |

\*  $p < 0.1$ , \*\*  $p < 0.05$ , \*\*\*  $p < 0.01$


<sup>1</sup> This table presents the 2SLS estimation of the regression (Equation ??) with school fixed effects. Column (1) displays the 2SLS estimation with the 15 question CES-D from the in-school survey. Column (2) displays the 2SLS estimation with the 19 question CES-D from the in-home survey in wave I. The control variables include sex, race, age, parental education, parental employment, parental occupation, and number of friends.

<sup>2</sup> Standard errors are clustered on the school level.

<sup>3</sup> Data source is the Add Health in-school and in-home surveys.


## Web.B Figures

Figure Online Appendix Figure A.1: Monotonicity Test



*Note:* This plot demonstrates the test for the monotonicity assumption in our instrumental variable approach. The y-axis shows the estimated effect of the instrument (friends-of-friends' mental unwellness) on the treatment (friends' mental unwellness) across different quantiles of the data. Monotonicity holds if the effect is consistently positive or negative across all quantiles. The dashed red line at  $y=0$  aids in assessing sign consistency. I also conduct a formal test of whether the slope of these quantile effects differs significantly from zero; a non-significant result ( $p > 0.05$ ) supports the monotonicity assumption. Violations of monotonicity may indicate the presence of 'defiers' in the sample, potentially biasing IV estimates.

Figure Online Appendix Figure A.2: Spillover Effect of Peers' Mental Unwellness on Long Run Outcomes




*Note:* See the notes in Figure ???. Each row on the y-axis is a separate regression with a different dependent variable. The x-axis is the coefficient of the mental unwellness of peers on the dependent variable. The dependent variables are (from bottom to the top): the probability of finishing high school in wave five, the probability of dropping out of high school in wave five, the probability of finishing college in wave five, the probability of being married in wave five, the number of times married in wave five (standardized), the probability of ever getting arrested in wave four, the probability of hurting others while on the influence of drugs in wave four, the probability of hurting others while on the influence of alcohol in wave four the probability of working full time in wave five, the probability of working full time in wave four, the number of weekly hours worked in wave five (standardized), the number of weekly hours worked in wave four (standardized), the probability of being employed in wave five, the probability of being employed in wave four, log hourly earnings in wave three, and log total personal earnings in wave four.

*Source:* The data source is the Add Health in-school and in-home surveys.




Figure Online Appendix Figure A.3: Spillover Effect of Peers' Mental Unwellness on Own Unwellness Health, Academic and Behavioral Outcomes




This plot presents the 2SLS estimation of the regression (equation ??) with school fixed effects on mental unwellness, academic, and behavioral outcomes. Additionally, separate 2SLS estimations of equation (??) are provided for different samples. In panel (A), the 2SLS estimation is presented for own mental unwellness. Panel (B) shows the 2SLS estimation for GPA. Panel (C) displays the 2SLS estimation for skipping school without an excuse. The control variables in the analysis include sex, race, age, parental education, parental employment, parental occupation, and number of friends. Standard errors are clustered at the school level. The data source is the Add Health in-school survey.

Figure Online Appendix Figure A.4: Spillover Effect of Peers' Mental Unwellness on Health Outcomes


(a) Poor Health



(b) Feel Sick




(c) Feel Tired




See the notes in Figure [Online Appendix Figure A.3](#). I have conducted separate 2SLS estimations for different samples. In panel (A), the 2SLS estimation is presented for the effect of equation (??) on self-reported poor health. Panel (B) provides the 2SLS estimation for the impact of equation (??) on feeling sick. In panel (C), the 2SLS estimation pertains to the effect of equation (??) on feeling tired. The control variables used in the analysis include sex, race, age, parental education, parental employment, parental occupation, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-school survey.

Figure Online Appendix Figure A.5: Spillover Effect of Peers' Mental Unwellness on Health Outcomes

(a) Never Seen a Therapist




(b) Miss School due to Health



15


See the notes in Figure Online Appendix Figure A.3. I have conducted separate 2SLS estimations for different samples. Panel (A) shows the 2SLS estimation for the influence of equation (??) on never seeing a therapist. Finally, in panel (B), the 2SLS estimation is presented for the relationship of equation (??) with missing school due to health reasons. The control variables used in the analysis include sex, race, age, parental education, parental employment, parental occupation, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-school survey.

Figure Online Appendix Figure A.6: Spillover Effect of Peers' Mental Unwellness on Behavioral Outcomes




See the notes in Figure [Online Appendix Figure A.3](#). In panel (A) demonstrates the 2SLS estimation for smoking, while Panel (B) covers intoxication. The control variables in the analysis include sex, race, age, parental education, parental employment, parental occupation, and number of friends. Standard errors are clustered at the school level. The data source is the Add Health in-school survey.

Figure Online Appendix Figure A.7: Spillover Effect of Peers' Mental Unwellness on Alcohol Consumption



See the notes in Figure [Online Appendix Figure A.3](#). All dependent variables are normalized to have a mean of zero and a standard deviation of one. Panel (A) the 2SLS estimation is presented for the frequency of alcohol use in wave 4. In panel (B), the 2SLS estimation is presented for the frequency of drug use in wave 3. For the analysis using in-home survey wave 3, control variables include sex, race, age, parental education, parental employment, parental health, and number of friends. For the analysis using in-home survey waves 4, control variables include sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-home survey.


Figure Online Appendix Figure A.8: Spillover Effect of Peers' Mental Unwellness on Socialization




See the notes in Figure [Online Appendix Figure A.3](#). All dependent variables are normalized to have a mean of zero and a standard deviation of one. In panel (A), the 2SLS estimation is presented for the number of close friends in wave 4. Panel (B) shows the 2SLS estimation is presented for the frequency a person hangs out with their friends in wave 5. For the analysis using in-home survey waves 4 and 5, control variables include sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-home survey.

Figure Online Appendix Figure A.9: Spillover Effect of Peers' Mental Unwellness on Drug Consumption


(a) Frequency of Marijuana Use (Wave 1)



(b) Frequency of Marijuana Use (Wave 3)




(c) Frequency of Marijuana Use (Wave 4)



See the notes in Figure [Online Appendix Figure A.3](#). All dependent variables are normalized to have a mean of zero and a standard deviation of one. Panels (A), (B), and (C) present the 2SLS estimation is presented for the frequency of marijuana use during the last 30 days in waves 1, 3, and 4, respectively. For the analysis using in-home survey waves 1 to 3, control variables include sex, race, age, parental education, parental employment, parental health, and number of friends. For the analysis using in-home survey waves 4 and 5, control variables include sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-home survey.


Figure Online Appendix Figure A.10: Spillover Effect of Peers' Mental Unwellness on Sexual Behavior



See the notes in Figure [Online Appendix Figure A.3](#). Panel (A) provides the 2SLS estimation is presented for having sex (wave 1). Panel (B) shows the 2SLS estimation is presented for intimate touching (wave 1). For the analysis using in-home survey wave 1, control variables include sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-home survey.




Figure Online Appendix Figure A.11: Spillover Effect of Peers' Mental Unwellness on BMI During Adolescents




See the notes in Figure Online Appendix Figure A.3. All the dependent variables are standardized with a mean of zero and a standard deviation of one. In panel (A), the 2SLS estimation is presented for BMI in wave 1. In panel (B), the 2SLS estimation is presented for BMI in wave 2. For the analysis that uses the in-home survey waves 1 to 3, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.

Figure Online Appendix Figure A.12: Spillover Effect of Peers' Mental Unwellness on BMI During Adulthood




See the notes in Figure [Online Appendix Figure A.3](#). All the dependent variables are standardized with a mean of zero and a standard deviation of one. In panel (A), the 2SLS estimation is presented for BMI in wave 3. In panel (B), the 2SLS estimation is presented for BMI in wave 4. In panel (C), the 2SLS estimation is presented for BMI in wave 5. For the analysis that uses the in-home survey wave 3, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. For the analysis that uses the in-home survey waves 4 and 5, I control for sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.

Figure Online Appendix Figure A.13: Spillover Effect of Mental Unwellness on Own Mental Unwellness During Adolescents




See the notes in Figure [Online Appendix Figure A.3](#). All the dependent variables are standardized with a mean of zero and a standard deviation of one. In panel (A), I provide the 2SLS estimation is presented for mental unwellness score in wave 1. In panel (B), I provide the 2SLS estimation is presented for mental unwellness score in wave 2. For the analysis that uses the in-home survey waves 1 to 3, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.

Figure Online Appendix Figure A.14: Spillover Effect of Mental Unwellness on Own Mental Unwellness During Adulthood



See the notes in Figure [Online Appendix Figure A.3](#). All the dependent variables are standardized with a mean of zero and a standard deviation of one. In panel (A), I provide the 2SLS estimation is presented for mental unwellness score in wave 3. In panel (B), I provide the 2SLS estimation is presented for mental unwellness score in wave 4. In panel (C), I provide the 2SLS estimation is presented for mental unwellness score in wave 5. For the analysis that uses the in-home survey wave 3, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. For the analysis that uses the in-home survey waves 4 and 5, I control for sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.


Figure Online Appendix Figure A.15: Spillover Effect of Mental Unwellness on Welfare Programs take-up




See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), I provide the 2SLS estimation is presented for receiving income from disability, unemployment, and social security benefits in wave 3. In panel (B), I provide the 2SLS estimation is presented for receiving food stamps in wave 3. In panel (C), I provide the 2SLS estimation is presented for receiving welfare payments in wave 3. For the analysis that uses the in-home survey wave 3, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. For the analysis that uses the in-home survey waves 4 and 5, I control for sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.

Figure Online Appendix Figure A.16: Spillover Effect of Mental Unwellness on Earnings

(a) Hourly Earnings (w3)




(b) Log Total Personal Earnings (w4)



See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), I provide the 2SLS estimation is presented for hourly earnings in wave 3. In panel (B), I provide the 2SLS estimation is presented for log total personal earnings in wave 4. For the analysis, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.


Figure Online Appendix Figure A.17: Spillover Effect of Mental Unwellness on Weekly Hours Worked



See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), I provide the 2SLS estimation is presented for weekly hours worked in wave 4. In panel (B), I provide the 2SLS estimation is presented for weekly hours worked in wave 5. For the analysis, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.


Figure Online Appendix Figure A.18: Spillover Effect of Mental Unwellness on Full Time Work

(a) Worked Full Time (w4)



Averages: Full sample 0.630, Women 0.630, Men 0.630,  
Above Median Num Frnds 0.640, Below Median Num Frnds 0.630.

(b) Worked Full Time (w5)




Averages: Full sample 0.680, Women 0.700, Men 0.660,  
Above Median Num Frnds 0.690, Below Median Num Frnds 0.680.

See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), I provide the 2SLS estimation is presented for working full time in wave 4. In panel (B), I provide the 2SLS estimation is presented for working full time in wave 5. For the analysis, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.




Figure Online Appendix Figure A.19: Spillover Effect of Mental Unwellness on Employment Status




See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), I provide the 2SLS estimation is presented for employment status in wave 4. In panel (B), I provide the 2SLS estimation is presented for employment status in wave 5. For the analysis, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.

Figure Online Appendix Figure A.20: Spillover Effect of Peers' Mental Unwellness on Schooling


(a) High School



(b) High School Dropout




(c) Finish College



See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), the 2SLS estimation is presented for finishing high school. In panel (B), the 2SLS estimation is presented for dropping out of high school. In panel (C), the 2SLS estimation is presented for finishing college. The control variables include sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey. The dependent variables in panels A to C are constructed from wave 5.


Figure Online Appendix Figure A.21: Spillover Effect of Peers' Mental Unwellness on Marriage

(a) Marital Status



Averages: Full sample 0.410, Women 0.470, Men 0.360,  
Above Median Num Frnds 0.430, Below Median Num Frnds 0.410.


(b) Times Married



Averages: Full sample 0.540 (0.580), Women 0.580 (0.590), Men 0.490 (0.570),  
Above Median Num Frnds 0.550 (0.580), Below Median Num Frnds 0.530 (0.580).


See the notes in Figure [Online Appendix Figure A.3](#). The dependent variable times married is normalized with a mean of zero and standard deviation of one. In panel (A), the 2SLS estimation is presented for being married in wave 5. In panel (B), the 2SLS estimation is presented for times married. The control variables include sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey. The dependent times married dependent variable is constructed using wave 4 data, while marital status is constructed using data from wave 5.

Figure Online Appendix Figure A.22: Spillover Effect of Peers' Mental Unwellness on Criminal Behavior




See the notes in Figure [Online Appendix Figure A.3](#). Panel (A) presents the 2SLS estimation is presented for if they have ever been arrested (wave 4). Panels (B) and (C) provide the 2SLS estimation is presented for if they ever hurt others while drunk or high—including unprotected sex (wave 4). For the analysis using in-home survey waves 1 to 3, control variables include sex, race, age, parental education, parental employment, parental health, and number of friends. For the analysis using in-home survey waves 4 and 5, control variables include sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-home survey.

Figure Online Appendix Figure A.23: Spillover Effect of Peers' Mental Unwellness on Sexually Risky Behavior




See the notes in Figure [Online Appendix Figure A.3](#). Panel (A) shows the 2SLS estimation is presented for having multiple sexual partners (wave 4). Panel (B) presents the 2SLS estimation is presented for if they ever paid for sex (wave 4). For the analysis using in-home survey waves 4, control variables include sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered at the school level. The data source for this study is the Add Health in-home survey.

Figure Online Appendix Figure A.24: Spillover Effect of Mental Unwellness on Smoking Cigarettes During Adolescents



See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), I provide the 2SLS estimation is presented for smoking cigarettes in wave 1. In panel (B), I provide the 2SLS estimation is presented for smoking cigarettes in wave 2. For the analysis that uses the in-home survey waves 1 to 3, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.

Figure Online Appendix Figure A.25: Spillover Effect of Mental Unwellness on Smoking Cigarettes During Adulthood



See the notes in Figure [Online Appendix Figure A.3](#). In panel (A), I provide the 2SLS estimation is presented for smoking cigarettes in wave 3. In panel (B), I provide the 2SLS estimation is presented for smoking cigarettes in wave 4. In panel (C), I provide the 2SLS estimation is presented for smoking cigarettes in wave 5. For the analysis that uses the in-home survey wave 3, I control for sex, race, age, parental education, parental employment, parental health, and number of friends. For the analysis that uses the in-home survey waves 4 and 5, I control for sex, race, age, educational attainment, parental education, parental employment, parental health, and number of friends. Standard errors are clustered on the school level. Data source is the Add Health in-home survey.